

LABIAPLASTY

MAKING AN INFORMED DECISION

GENERAL INFORMATION

Labiaplasty is a surgical procedure to remove excess redundant labia, which cause unsightly contour lines and physical discomfort. Some women report pinching or chafing when sitting or walking and the redundant skin can hinder sexual activity.

ALTERNATIVE TREATMENTS

Alternative forms of management consist of not treating the labia.

RISKS OF LABIAPLASTY SURGERY

Every surgical procedure involves a certain amount of risk and it is important that you understand these risks and the possible complications associated with them. In addition, every procedure has limitations. An individual's choice to undergo a surgical procedure is based on the comparison of the risk to potential benefit. Although the majority of patients do not experience these complications, you should discuss each of them with Dr Linsell to make sure you completely understand all possible consequences of a Labiaplasty.

Bleeding- It is possible, though unusual, to experience a bleeding episode during or after surgery. Should post-operative bleeding occur, it may require an emergency treatment to drain the accumulated blood or blood transfusion. Intra-operative blood transfusions may be required. Do not take any aspirin or anti-inflammatory medications for ten days before surgery, as this may increase the risk of bleeding. Non-prescription "herbs" and dietary supplements can increase the risk of surgical bleeding. Hematoma can occur at any time following injury. If blood transfusions are needed to treat blood loss, there is a risk of blood related infections such as hepatitis and the HIV (AIDS). Heparin medications that are used to prevent blood clots in veins can produce bleeding and decreased blood platelets.

Infection - Infection is unusual after surgery as you are routinely given antibiotics during and for 10 days after your procedure. Infection may contribute to breakdown of your wound and delayed healing. This will always heal yet may take longer than expected. Should an infection occur, treatment including antibiotics, hospitalization, or additional surgery may be necessary.

Change in Skin Sensation- It is common to experience diminished (or loss) of skin sensation in areas that have had surgery. Itching, tenderness, or exaggerated responses to hot or cold temperatures may occur. Usually this resolves during healing, but in rare situations it may be chronic. Increased sensation, diminished sensation (or complete loss of skin sensation) may not totally resolve after a Labiaplasty which can change the sensation of the labia during intercourse.

Skin Contour Irregularities- Contour and shape irregularities and depressions may occur after Labiaplasty. Visible and palpable wrinkling of skin can occur. Residual skin irregularities at the ends of the incisions or "dog ears" are always a possibility as is skin pleating when there is excessive redundant skin. This may improve with time, or it can be surgically corrected with a second procedure.

PLEASE INITIAL HERE TO INDICATE YOU HAVE READ AND UNDERSTAND PAGE 1 OF THIS CONSENT FORM:

Major Wound Separation- Wounds may separate after surgery. Should this occur, additional treatment including surgery may be necessary.

Skin Discoloration / Swelling- Bruising and swelling normally occurs following Labiaplasty. The skin in or near the surgical site can appear either lighter or darker than surrounding skin. Although uncommon, swelling and skin discoloration may persist for long periods of time and, in rare situations, may be permanent.

Sutures- Dr Linsell uses dissolving sutures so there are no sutures to be removed after your operation. Occasionally, you may notice these sutures after your surgery. Sutures may spontaneously poke through the skin, become visible or produce irritation that requires removal.

Damage to Deeper Structures- There is the potential for injury to deeper structures including nerves, blood vessels, and muscles, during any surgical procedure. The potential for this to occur varies according to the type of procedure being performed. Injury to deeper structures may be temporary or permanent.

Scarring- All surgery leaves scars, some more visible than others. Although good wound healing after a surgical procedure is expected, abnormal scars may occur within the skin and deeper tissues. The larger the labia are to start with, the higher the likelihood of a poor scar. Scars may be unattractive, wide and of different colour than surrounding skin. Scar appearance may also vary within the same scar, exhibit contour variations or "bunching" due to the amount of excess skin. Scars may be asymmetrical (appear different between right and left side of the body). There is the possibility of visible marks in the skin from sutures. In some cases scars may require surgical revision or treatment.

Surgical Anaesthesia- Both local and general anaesthesia involve risk. There is the possibility of complications, injury, and even death from all forms of surgical anaesthesia or sedation

Asymmetry- Labiaplasty may result in an asymmetry of the body appearance. Factors such as skin tone, fatty deposits, skeletal prominence, and muscle tone may contribute to normal asymmetry in body features. Additional surgery may be necessary to attempt to improve asymmetry.

Allergic Reactions- In rare cases, local allergies to tape, suture material and glues, blood products, topical preparations or injected agents have been reported. Serious systemic reactions including shock (anaphylaxis) may occur to drugs used during surgery and prescription medications. Allergic reactions may require additional treatment.

Delayed Healing- Wound disruption or delayed wound healing is possible. Some areas of the labia may not heal normally and may take a long time to heal. Some areas of skin or tissue may die. This may require frequent dressing changes or further surgery to remove the non-healed tissue. **Smokers have a greater risk of skin loss and wound healing complications.**

Shock- In rare circumstances, your surgical procedure can cause severe trauma, particularly when multiple or extensive procedures are performed. Although serious complications are infrequent, infections or excessive fluid loss can lead to severe illness and even death. If surgical shock occurs, hospitalization and additional treatment would be necessary.

Persistent Swelling (Lymphedema)- Though rare, persistent swelling in the labia can occur following Labiaplasty.

PLEASE INITIAL HERE TO INDICATE YOU HAVE READ AND UNDERSTAND PAGE 2 OF THIS CONSENT FORM:

Pain- You will experience pain after your surgery. Pain of varying intensity and duration may occur and persist after Labiaplasty. Dr Linsell has a comprehensive program to minimise your pain and you are advised to follow this religiously. Chronic pain may occur very infrequently from nerves becoming trapped in scar tissue after Labiaplasty, which may require treatment with drugs or further surgery.

Unsatisfactory Result- Although good results are expected, there is no guarantee or warranty expressed or implied, on the results that may be obtained. You may be disappointed with the results of Labiaplasty surgery. This would include risks such as asymmetry, unsatisfactory or highly visible surgical scar location, unacceptable visible deformities, bunching and rippling in the skin near the suture lines or at the ends of the incisions (dog ears), poor healing, wound disruption, and loss of sensation. It may not be possible to correct or improve the effects of surgical scars. In some situations, it may not be possible to achieve optimal results with a single surgical procedure. Additional surgery may be required to improve results.

Deep Venous Thrombosis, Cardiac and Pulmonary Complications- Surgery, especially longer procedures, may be associated with the formation of, or increase in, blood clots in the venous system. Pulmonary complications may occur secondarily to both blood clots (pulmonary emboli), fat deposits (fat emboli) or partial collapse of the lungs after general anaesthesia. Pulmonary and fat emboli can be life- threatening or fatal in some circumstances. Air travel, inactivity and other conditions may increase the incidence of blood clots traveling to the lungs causing a major blood clot that may result in death. It is important to discuss with Dr Linsell any past history of blood clots, swollen legs or the use of oestrogen or birth control pills that may contribute to this condition. Cardiac complications are a risk with any surgery and anaesthesia, even in patients without symptoms. Heart attack or stroke could occur due to strain on the heart. Should any of these complications occur, you may require hospitalization and additional treatment. If you experience shortness of breath, chest pains, or unusual heart beats, seek medical attention immediately.

ADDITIONAL ADVICE

Long-Term Results- Subsequent alterations in the appearance of your body may occur as the result of aging, sun exposure, weight loss, weight gain, pregnancy, menopause or other circumstances not related to your surgery.

Body-Piercing Procedures- Individuals who currently wear body-piercing jewellery or are seeking to undergo body-piercing procedures must consider the possibility that an infection could develop anytime following this procedure. Treatment including antibiotics, hospitalisation or additional surgery may be necessary.

Female Patient Information- It is important to inform your plastic surgeon if you use birth control pills, oestrogen replacement, or if you suspect you may be pregnant. Many medications including antibiotics may neutralize the preventive effect of birth control pills, allowing for conception and pregnancy.

Intimate Relations After Surgery- Surgery involves coagulating of blood vessels and increased activity of any kind may open these vessels leading to a bleed, or haematoma. Increased activity that increased your pulse or heart rate may cause additional bruising, swelling, and the need for return to surgery and control bleeding. It is wise to refrain from sexual activity for about three weeks after your operation.

PLEASE INITIAL HERE TO INDICATE YOU HAVE READ AND UNDERSTAND PAGE 3 OF THIS CONSENT FORM:

Medications- There are many adverse reactions that occur as the result of taking over-the-counter, herbal, and/or prescription medications. Be sure to check with Dr Linsell about any drug interactions that may exist with medications that you are already taking. If you have an adverse reaction, stop the drugs immediately and call Dr Linsell for further instructions. If the reaction is severe, go immediately to the nearest emergency room. When taking the prescribed pain medications after surgery, realize that they can affect your thought process and coordination. Do not drive, do not operate complex equipment, do not make any important decisions and do not drink any alcohol while taking these medications. Be sure to take your prescribed medication only as directed.

Mental Health Disorders and Elective Surgery- It is important that all patients seeking to undergo elective surgery have realistic expectations that focus on improvement rather than perfection. Complications or less than satisfactory results are sometimes unavoidable, may require additional surgery and often are stressful. Please openly discuss with Dr Linsell, prior to surgery, any history that you may have of significant emotional depression or mental health disorders. Although many individuals may benefit psychologically from the results of elective surgery, effects on mental health cannot be accurately predicted.

ADDITIONAL SURGERY NECESSARY (RE-OPERATIONS)

There are many variable conditions that may influence the long-term result of surgery. Should complications occur, additional surgery or other treatments may be necessary. Secondary surgery may be necessary to obtain optimal results. Even though risks and complications occur infrequently, the risks cited are particularly associated with a Labiaplasty. The practice of medicine and surgery is not an exact science. Although good results are expected, there is no guarantee or warranty expressed or implied, on the results that may be obtained. In some situations, it may not be possible to achieve optimal results with a single surgical procedure.

PATIENT COMPLIANCE

Please follow Dr Linsell's instructions carefully. This is essential for the success of your outcome. It is important that the surgical incisions are not subjected to excessive force, swelling, abrasion, or motion during the time of healing. Personal and vocational activity needs to be restricted. Successful post-operative function depends on both surgery and subsequent care. Physical activity that increases your pulse or heart rate may cause bruising, swelling, fluid accumulation and the need for return to surgery. It is important that you participate in follow-up care, return for aftercare, and promote your recovery after surgery.

FINANCIAL RESPONSIBILITIES

The cost of surgery involves several charges for the services provided. The total includes fees charged by Dr Linsell, the anaesthetist, the surgical assistant, laboratory tests, and possible hospital charges. If your surgery attracts a Medicare item number, you may receive some funds back from Medicare and where applicable, your health fund. The fees charged for this procedure do not include any potential future costs for additional procedures that you elect to have or require in order to revise, optimize, or complete your outcome. Additional costs may occur should complications develop from the surgery. Whilst Dr Linsell will generally perform revisional surgery within the first 12 months for no out of pocket cost for his fee, there may be charges for the anaesthetist, assistant and hospital.

PLEASE INITIAL HERE TO INDICATE YOU HAVE READ AND UNDERSTAND PAGE 4 OF THIS CONSENT FORM:

HEALTH INSURANCE

There is a Medicare item number for Labiaplasty (35534), however most people do not qualify for this item number. If you do, most health insurance companies cover some of the costs for the procedure. Please carefully review your health insurance subscriber-information or contact your insurance company for a detailed explanation of their policies for covering Labiaplasty procedures. **Some health insurance plans exclude coverage for secondary or revisionary surgery.**

DISCLAIMER

Informed-consent documents are used to communicate information about the proposed surgical treatment of a disease or condition along with disclosure of risks and alternative forms of treatment(s), including no surgery. The informed-consent process attempts to define principles of risk disclosure that should generally meet the needs of most patients in most circumstances.

However, informed-consent documents should not be considered all-inclusive in defining other methods of care and risks encountered. Dr Linsell may provide you with additional or different information which is based on all the facts in your particular case and the current state of medical knowledge.

Informed-consent documents are not intended to define or serve as the standard of medical care. Standards of medical care are determined on the basis of all of the facts involved in an individual case and are subject to change as scientific knowledge and technology advance and as practice patterns evolve.

NO SURGERY IS GUARANTEED.

It is important that you read the above information carefully and have all of your questions answered before signing the consent on the next page.

PLEASE INITIAL HERE TO INDICATE YOU HAVE READ AND UNDERSTAND PAGE 5 OF THIS CONSENT FORM:

CONSENT FORM – LABIOPLASTY

The following form acknowledges my willingness to undergo the procedure of Labiaplasty performed by Dr Malcolm Linsell.

Initial:

I acknowledge that I have consulted with Dr Linsell and I have a good understanding of the operation and the expected outcomes.

Initial:

I acknowledge the cost of the procedure has been explained in detail and that there will be fees for surgeon, anaesthetist, assistant surgeon and the hospital.

Initial:

I acknowledge that this is real surgery and any surgery carries with it the risk of complications.

Initial:

I acknowledge Dr Linsell has explained the most common complications that can arise from this procedure.

Initial:

I acknowledge and accept that some complications can interfere with my expected outcome.

Initial:

I acknowledge I have been given ample time and opportunity to ask Dr Linsell questions about the procedure, my expected outcome and the possible complications.

Initial:

I acknowledge that photographs of my labia will be taken before, during and after the operation.

Initial:

I acknowledge and agree to the use of my photographs to follow my own progress and (without being identified) to share with other patients to help them with their decision-making.

Initial:

I agree to the potential use of my photographs (without being identified) in medical conferences and publications.

Initial:

I acknowledge blood products or derivatives are occasionally necessary to administer during and after the operation.

Initial:

I acknowledge that in the event of one of the hospital staff coming into direct contact with my blood, blood samples will be taken from the staff member and myself to test for infections such as Hepatitis and HIV.

Initial:

I acknowledge a revision surgical procedure is occasionally necessary for up to 12 months after the procedure. If required, Dr Linsell will do this for a small out of pocket cost for his surgical fee, however there may be fees for anaesthetist, assistant surgeon and hospital.

Initial:

My signature below confirms I have read and understand the information provided and hereby consent to the procedure.

Print name:

Signed:

Date:

Dr Malcolm D Linsell

Signed:

Date: